

Centrally symmetric generators in toric Fano varieties*

Cinzia Casagrande[†]

22nd October 2002

Abstract. We give a structure theorem for n -dimensional smooth toric Fano varieties whose associated polytope has “many” pairs of centrally symmetric vertices.

Introduction. Smooth toric Fano varieties, together with their equivariant birational contractions, have been intensively studied in recent years: see [WW82, Bat82, Bat99, Sat00, Bon00, Cas01b, Cas01c, Sat02]. In some cases, the toric case has been a testing ground for results concerning general smooth Fano varieties, see for instance [BCW02, BCDD02]. Despite the strong combinatorial properties of fans of toric Fano varieties, these varieties are classified only in dimension less or equal to 4 (see [WW82, Bat82] for dimension 3, [Bat99] and [Sat00], example 4.7, for dimension 4).

Let X be a smooth, complete toric variety of dimension n , $N = \text{Hom}_{\mathbb{Z}}(\mathbb{C}^*, T)$, $N_{\mathbb{Q}} = N \otimes_{\mathbb{Z}} \mathbb{Q}$ and Σ_X the fan of X in $N_{\mathbb{Q}}$. We denote by $\langle x_1, \dots, x_h \rangle$ the convex cone spanned in $N_{\mathbb{Q}}$ by $x_1, \dots, x_h \in N$.

Definition. The set of *generators* of Σ_X is defined in the following way:

$$G(\Sigma_X) = \{x \in N \mid \langle x \rangle \in \Sigma_X \text{ and } x \text{ is primitive in } \langle x \rangle \cap N\}.$$

Suppose that X is Fano. Then the generators $G(\Sigma_X)$ are the vertices of a simplicial convex polytope $\mathcal{P}_X \subset N_{\mathbb{Q}}$, and every cone in Σ_X is the cone over a face of $\mathcal{P}_X$. Hence the set of generators $G(\Sigma_X)$ determines the fan Σ_X (and X). This means that X is the only Fano variety in the set

$$\begin{aligned} \mathcal{B}_X &= \{X' \text{ toric, smooth, complete} \mid \exists \varphi: X' \dashrightarrow X \text{ equivariant, birational, isomorphism} \\ &\quad \text{in codimension 1}\} \\ &= \{X' \text{ toric, smooth, complete} \mid G(\Sigma_{X'}) = G(\Sigma_X)\} \end{aligned}$$

and any such X' determines X . It is an interesting question whether there exists some X' in $\mathcal{B}_X$ with a particularly simple structure. More precisely, we think of a toric bundle structure (see section 2), with fiber and basis some lower dimensional toric Fano variety. This would give an inductive way to classify these varieties. Our result is the following:

*2000 Mathematics Subject Classification: 14J45, 14M25, 52B20.

[†]Dipartimento di Matematica, Università di Roma “La Sapienza”, piazzale Aldo Moro 5, 00185 Roma, e-mail: ccasagra@mat.uniroma1.it

Theorem (theorem 5 and proposition 7). *Let X be a smooth toric Fano variety. Suppose that $G(\Sigma_X)$ contains $r > 0$ pairs of centrally symmetric generators $\{x, -x\}$, and consider the linear subspace H of $N_{\mathbb{Q}}$ spanned by all such pairs.*

- *If $\dim H < r$, then X is a toric bundle over a lower dimensional toric Fano variety, with fiber a product of Del Pezzo varieties (see definition on page 4).*
- *If H contains more than $2 \dim H$ generators, then there exists X' in $\mathcal{B}_X$ which is a toric bundle over a lower dimensional toric Fano variety, with fiber a product of Del Pezzo and pseudo Del Pezzo varieties (see definitions on page 4).*
- *If $\dim H = n - 1$, then there exists X' in $\mathcal{B}_X$ which is a toric bundle over $\mathbb{P}^1$, with fiber a product of Del Pezzo varieties, pseudo Del Pezzo varieties and a power of $\mathbb{P}^1$.*
- *If $\dim H = n$, then X itself is a product of Del Pezzo varieties, pseudo Del Pezzo varieties and a power of $\mathbb{P}^1$.*

All these results give a structure theorem for some classes of toric Fano varieties, in any dimension. The last case ($\dim H = n$) generalizes a previous result by G. Ewald [Ewa88], who obtains the same statement under the stronger hypothesis that there exists an n -dimensional cone $\sigma \in \Sigma_X$ such that $-\sigma \in \Sigma_X$.

The structure of the paper is as follows: in section 1 we recall the language of primitive relations and some results of toric Mori theory; in section 2 we recall properties of toric bundles; in section 3 we introduce Del Pezzo and pseudo Del Pezzo varieties and state and prove our results.

1. Primitive relations and 1-cycles. We collect here some basic notions and results. We refer to [Oda88, Ful93] for properties of toric varieties, to [Bat99] for primitive collections and primitive relations, to [Rei83] and to the survey paper [Wiś02] for toric Mori theory.

Let X be a smooth, complete toric variety.

Definition. A set $P \subseteq G(\Sigma_X)$ is a *primitive collection* if $\langle P \rangle \notin \Sigma_X$ and $\langle P \setminus \{x\} \rangle \in \Sigma_X$ for all $x \in P$.

If $P = \{x_1, \dots, x_h\}$ is a primitive collection, consider the minimal cone $\sigma \in \Sigma_X$ containing the point $x_1 + \dots + x_h$. If $\sigma = \langle y_1, \dots, y_k \rangle$, we get a relation

$$x_1 + \dots + x_h - (a_1 y_1 + \dots + a_k y_k) = 0$$

with $a_i \in \mathbb{Z}_{>0}$. This is the *primitive relation* $r(P)$ associated to P , and the *degree* of P is $\deg P = h - \sum_i a_i$.

Let $\mathcal{A}_1(X)$ be the group of algebraic 1-cycles on X modulo numerical equivalence, $\mathcal{N}_1(X) = \mathcal{A}_1(X) \otimes_{\mathbb{Z}} \mathbb{Q}$, and $\text{NE}(X) \subset \mathcal{N}_1(X)$ the convex cone generated by classes of effective curves; we recall that $\text{NE}(X)$ is closed and polyhedral.

For every $r \in \{0, \dots, n\}$ there is a bijection between the set of r -dimensional cones in Σ_X and the set of codimension r orbits of T in X . For $r = 1$, this gives a bijection between $G(\Sigma_X)$ and the set of codimension 1 orbits of T . We denote by $V(\sigma)$ (respectively $V(x)$)

the closure of the orbit corresponding to $\sigma \in \Sigma_X$ (respectively $x \in G(\Sigma_X)$); this is an irreducible invariant subvariety.

We recall that there is an exact sequence:

$$0 \longrightarrow \mathcal{A}_1(X) \longrightarrow \mathbb{Z}^{G(\Sigma_X)} \longrightarrow N \longrightarrow 0$$

where the map $\mathcal{A}_1(X) \rightarrow \mathbb{Z}^{G(\Sigma_X)}$ is given by $\gamma \mapsto \{\gamma \cdot V(x)\}_{x \in G(\Sigma_X)}$. Hence the group $\mathcal{A}_1(X)$ is canonically isomorphic to the lattice of integral relations among the elements of $G(\Sigma_X)$: a relation $\sum_x a_x x = 0$ corresponds to a numerical class γ that has intersection a_x with each divisor $V(x)$. Moreover, we have $(-K_X) \cdot \gamma = \sum_x a_x$. In particular, every primitive relation $r(P)$ defines a numerical class in $\mathcal{A}_1(X)$, and $\deg P = (-K_X) \cdot r(P)$. In the sequel, we will identify numerical classes with their associated relations. Remark that from the exact sequence above we deduce $\rho_X = \#G(\Sigma_X) - n$, where ρ_X is the Picard number of X .

Lemma 1 ([Cas01a], lemma 1.4). *Let $\gamma \in \mathcal{A}_1(X)$ given by the relation*

$$a_1 x_1 + \cdots + a_h x_h - (b_1 y_1 + \cdots + b_k y_k) = 0$$

with $a_i, b_j \in \mathbb{Z}_{>0}$ for each i, j . If $\langle y_1, \dots, y_k \rangle \in \Sigma_X$, then $\gamma \in \text{NE}(X)$.

In particular, every primitive relation is an effective class: therefore, if X is Fano, every primitive collection has strictly positive degree.

Theorem 2. *Let X be a toric Fano variety and let $\gamma \in \text{NE}(X) \cap \mathcal{A}_1(X)$ be such that $(-K_X) \cdot \gamma = 1$. Then γ is extremal in $\text{NE}(X)$, the relation associated to γ is a primitive relation*

$$x_1 + \cdots + x_h - (a_1 y_1 + \cdots + a_k y_k) = 0,$$

and for every cone $\nu = \langle z_1, \dots, z_t \rangle$ such that $\{z_1, \dots, z_t\} \cap \{x_1, \dots, x_h, y_1, \dots, y_k\} = \emptyset$ and $\langle y_1, \dots, y_k \rangle + \nu \in \Sigma_X$, we have $\langle x_1, \dots, x_h, y_1, \dots, y_k \rangle + \nu \in \Sigma_X$ for all $i = 1, \dots, h$.

Proof. See [Cas01a], corollary 4.4 and [Rei83], theorem 2.4. ■

Lemma 3 ([Cas01c], lemma 2.2). *Let X be a toric Fano variety. Suppose that $\{x, -x\}$ is a primitive collection in Σ_X . Then for any other primitive collection P containing x , the associated primitive relation is*

$$r(P) : \quad x + y_1 + \cdots + y_h - (z_1 + \cdots + z_h) = 0;$$

moreover, $P' = \{-x, z_1, \dots, z_h\}$ is also a primitive collection, with relation

$$r(P') : \quad (-x) + z_1 + \cdots + z_h - (y_1 + \cdots + y_h) = 0.$$

Both P and P' have degree 1, their primitive relations are extremal, and $2h \leq n$.

2. Toric bundles. We recall the following standard result of toric geometry (see [Ewa96], theorem 6.7 in chapter VI, or [Ful93], section 2.4, exercise on fiber bundles):

Theorem 4. *Let X be a smooth, complete toric variety. Consider a linear subspace $H \subset N_{\mathbb{Q}}$ of dimension r and suppose that for every n -dimensional cone $\sigma \in \Sigma_X$, we have $\sigma = \eta + \tau$, with $\eta \in \Sigma_X$, $\eta \subset H$, $\dim \eta = r$ and $\tau \cap H = \{0\}$. Then:*

- *the set $\Sigma_F = \{\sigma \in \Sigma_X \mid \sigma \subset H\}$ is the fan of a smooth, complete, r -dimensional toric variety F ;*
- *consider a complementar $H^{\perp}$ of H in $N_{\mathbb{Q}}$ such that $N = (N \cap H) \oplus (N \cap H^{\perp})$, and let $\pi: N_{\mathbb{Q}} \rightarrow H^{\perp}$ be the projection. Then the set $\Sigma_Z = \{\pi(\sigma) \mid \sigma \in \Sigma_X\}$ is the fan of a complete, smooth, $(n - r)$ -dimensional toric variety Z ;*
- *the projection π induces an equivariant morphism $\bar{\pi}: X \rightarrow Z$ such that, for every affine invariant open subset $U \subset Z$, $\bar{\pi}^{-1}(U) \simeq F \times U$ as toric varieties over U .*

Hence, X is a toric bundle over Z with fiber F .

Under this description, for every $\eta \in \Sigma_X$ such that $\eta \subset H$ and $\dim \eta = r$, the subvariety $V(\eta)$ is an invariant section of the bundle; for every $\tau \in \Sigma_X$ such that $\tau \cap H = \{0\}$ and $\dim \tau = n - r$, the subvariety $V(\tau)$ is an invariant fiber.

Suppose that X is a toric bundle over Z with fiber F , as described in theorem 4. Then $G(\Sigma_F) = G(\Sigma_X) \cap H$ and the primitive collections of Σ_X contained in H are exactly the primitive collections of Σ_F . These primitive collections have the same primitive relations in Σ_X and in Σ_F .

Now set $\mathcal{G} = G(\Sigma_X) \setminus G(\Sigma_F)$. The projection $\pi: N_{\mathbb{Q}} \rightarrow H^{\perp}$ induces a bijection between $\mathcal{G}$ and $G(\Sigma_Z)$. Under this identification, the primitive collections of Σ_X not contained in H are exactly the primitive collections of Σ_Z . We denote by $\bar{x}$ the image of an element $x \in N$ under π . A primitive relation in Σ_Z of the form

$$\bar{z}_1 + \cdots + \bar{z}_h - (a_1 \bar{y}_1 + \cdots + a_k \bar{y}_k) = 0$$

will lift in Σ_X to a primitive relation of the form:

$$z_1 + \cdots + z_h - (a_1 y_1 + \cdots + a_k y_k + b_1 x_1 + \cdots + b_l x_l) = 0,$$

with $l \geq 0$ and $x_i \in G(\Sigma_F)$ for all $i = 1, \dots, l$.

The bundle is trivial, namely $X \simeq F \times Z$, if and only if all primitive relations in Σ_Z remain unchanged when lifted in Σ_X .

3. Del Pezzo varieties and centrally symmetric generators. Let $x_1, \dots, x_n$ be a basis of the lattice N and suppose that $n = 2r$ is even. Consider the following elements of N :

$$x_0 = -x_1 - \cdots - x_n, \quad y_i = -x_i \quad \text{for } i = 0, \dots, n.$$

Notation: I^s denotes a set of indexes of cardinality s .

Definition. The n -dimensional *Del Pezzo variety* V^n is the toric Fano variety described by the fan $\Sigma_{V^n} \subset N_{\mathbb{Q}}$ such that $G(\Sigma_{V^n}) = \{x_0, y_0, \dots, x_n, y_n\}$.

The n -dimensional *pseudo Del Pezzo variety* $\tilde{V}^n$ is the toric Fano variety described by the fan $\Sigma_{\tilde{V}^n} \subset N_{\mathbb{Q}}$ such that $G(\Sigma_{\tilde{V}^n}) = \{x_0, x_1, y_1, \dots, x_n, y_n\}$.

We have $\rho_{V^n} = n + 2$ and $\rho_{\tilde{V}^n} = n + 1$. In dimension 2, these varieties are the Del Pezzo surfaces S_3 and S_2 , obtained from $\mathbb{P}^2$ blowing-up respectively three or two fixed points. If the dimension is greater than two, the varieties V^n and $\tilde{V}^n$ do not admit any smooth equivariant blow-down; moreover, the blow-up along any invariant subvariety is never Fano. This behaviour is rather peculiar: in dimension 4, H. Sato ([Sat00], theorem 8.1) has shown that any toric Fano 4-fold different from V^4 and $\tilde{V}^4$ can be obtained from $\mathbb{P}^4$ by a sequence of smooth equivariant blow-ups and blow-downs in such a way that all the intermediate varieties are Fano.

The variety V^n can be geometrically described as follows. Consider the Cremona map $\mathbb{P}^n \dashrightarrow \mathbb{P}^n$. It is well known that this map factorizes as a sequence of smooth blow-ups and blow-downs in the following way: first blow-up $n + 1$ general points $p_1, \dots, p_{n+1}$ in $\mathbb{P}^n$. Then, for all $m = 2, \dots, n - 2$ (in this order), flip the strict transforms of all the $\mathbb{P}^{m-1}$ spanned by m of the points p_i . Finally, blow-down the strict transforms of all the $\mathbb{P}^{n-1}$ spanned by n of the points p_i . It is easy to see that this is a toric factorization, namely, all varieties involved are toric and the maps are equivariant. Now, V^n is the intermediate variety that you get if you perform the flips just till the dimension $m = r - 1$, where $r = n/2$.

Del Pezzo varieties were introduced by V. E. Voskresenskii and A. A. Klyachko [VK85], where they show (theorem 6) that *every toric Fano variety whose fan is centrally symmetric is a product of Del Pezzo varieties and a power of $\mathbb{P}^1$* . This result has been generalized by G. Ewald [Ewa88], who introduces pseudo Del Pezzo varieties and shows that *every toric Fano variety whose fan contains two centrally symmetric maximal cones, is a product of Del Pezzo varieties, pseudo Del Pezzo varieties and a power of $\mathbb{P}^1$* . We will see in proposition 7 that it is actually enough to ask that Σ_X has n linearly independent pairs of symmetric generators.

Explicitly, the fans of V^n and $\tilde{V}^n$ are:

$$\begin{aligned}\Sigma_{V^n} &= \{ \langle x_i, y_j \rangle_{i \in I^r, j \in J^r} \mid I^r, J^r \subset \{0, \dots, n\} \text{ disjoint} \}; \\ \Sigma_{\tilde{V}^n} &= \{ \langle x_0, x_i, y_j \rangle_{i \in I^{r-1}, j \in J^r}, \langle x_i, y_j \rangle_{i \in \tilde{I}^{r+s}, j \in \tilde{J}^{r-s}} \mid I^{r-1}, J^r \subset \{1, \dots, n\} \text{ disjoint}, \\ &\quad s \in \{0, \dots, r\} \text{ and } \tilde{I}^{r+s}, \tilde{J}^{r-s} \text{ a partition of } \{1, \dots, n\} \}.\end{aligned}$$

The primitive relations of Σ_{V^n} are $x_i + y_i = 0$ for $i = 0, \dots, n$ and

$$\sum_{i \in I^{r+1}} x_i - \left(\sum_{j \in J^r} y_j \right) = 0, \quad \sum_{i \in I^{r+1}} y_i - \left(\sum_{j \in J^r} x_j \right) = 0$$

for every partition I^{r+1}, J^r of $\{0, \dots, n\}$.

The primitive relations of $\Sigma_{\tilde{V}^n}$ are $x_i + y_i = 0$ for $i = 1, \dots, n$ and

$$x_0 + \sum_{i \in I^r} x_i - \left(\sum_{j \in J^r} y_j \right) = 0, \quad \sum_{j \in J^{r+1}} y_j - \left(x_0 + \sum_{i \in I^{r-1}} x_i \right) = 0$$

for all partitions I^r, J^r and I^{r-1}, J^{r+1} of $\{1, \dots, n\}$.

What makes Del Pezzo and pseudo Del pezzo varieties special among toric Fano varieties, is that they have lots of numerical classes of curves with anticanonical degree 1.

Theorem 5. *Let X be a toric Fano variety and let H be the linear subspace of $N_{\mathbb{Q}}$ spanned by all pairs of centrally symmetric generators in $G(\Sigma_X)$. Then there exists a decomposition*

$$H = H_1 \oplus \cdots \oplus H_a \oplus \tilde{H}_1 \oplus \cdots \oplus \tilde{H}_b \oplus K$$

with $a \geq 0$, $h_i = \dim H_i > 0$, $b \geq 0$, $\tilde{h}_i = \dim \tilde{H}_i > 0$ and $\dim K \geq 0$, such that:

- (i) $G(\Sigma_X) \cap H = \bigcup_{i=1}^a (G(\Sigma_X) \cap H_i) \cup \bigcup_{j=1}^b (G(\Sigma_X) \cap \tilde{H}_j) \cup (G(\Sigma_X) \cap K)$;
- (ii) $\#(G(\Sigma_X) \cap H_i) = 2(h_i + 1)$, $\#(G(\Sigma_X) \cap \tilde{H}_j) = 2\tilde{h}_j + 1$ and $\#(G(\Sigma_X) \cap K) = 2 \dim K$ for all $i = 1, \dots, a$ and $j = 1, \dots, b$;
- (iii) there exists a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & X' \\ \downarrow & & \downarrow \\ Z & \xrightarrow{\psi} & Z' \longrightarrow W \end{array}$$

where all varieties are smooth, projective and toric, Z and W are Fano, φ and ψ are equivariant birational maps which are isomorphisms in codimension 1, and the decomposition of H induces toric bundle structures:

- $X \rightarrow Z$ and $X' \rightarrow Z'$ with fiber $V^{h_1} \times \cdots \times V^{h_a}$,
- $Z' \rightarrow W$ with fiber $\tilde{V}^{\tilde{h}_1} \times \cdots \times \tilde{V}^{\tilde{h}_b}$,
- $X' \rightarrow W$ with fiber $V^{h_1} \times \cdots \times V^{h_a} \times \tilde{V}^{\tilde{h}_1} \times \cdots \times \tilde{V}^{\tilde{h}_b}$.

Remarks:

- $X' \in \mathcal{B}_X$.
- in particular, we will see in the proof (step 7) that φ is an isomorphism over some invariant section of $X' \rightarrow W$, hence there exists an invariant subvariety $V \simeq W$ of X : V is Fano, $\dim V = n - \sum_i h_i - \sum_j \tilde{h}_j$ and $\rho_V = \rho_X - \sum_i (h_i + 2) - \sum_j (\tilde{h}_j + 1)$ (where ρ_V and ρ_X are the Picard number of V and X respectively).
- when H_i or $\tilde{H}_i$ has dimension 2, we recover [Cas01c], 2.4.3 and 2.4.2a respectively.
- for all $i = 1, \dots, a$, $G(\Sigma_X) \cap H_i$ consists of $h_i + 1$ pairs of symmetric generators $\{x_l, y_l\}$, h_i by h_i linearly independent, such that $x_1 + \cdots + x_{h_i+1} = 0$.
- for all $j = 1, \dots, b$, $G(\Sigma_X) \cap \tilde{H}_j$ consists of $\tilde{h}_j$ linearly independent pairs of symmetric generators $\{x_l, y_l\}$, plus another generator $x_0 = x_1 + \cdots + x_{\tilde{h}_j}$.
- $G(\Sigma_X) \cap K$ consists of $\dim K$ linearly independent pairs of symmetric generators.

For the proof of theorem 5, we need the following lemma:

Lemma 6. Suppose that Σ_X contains m pairs of centrally symmetric generators $x_i + y_i = 0$ for $i = 1, \dots, m$. Let $\tau = \langle z_1, \dots, z_s \rangle \in \Sigma_X$ such that $\{z_1, \dots, z_s, x_1, \dots, x_m\}$ are linearly independent. Then there exists a partition I, J of $\{1, \dots, m\}$ such that $\tau + \langle x_i, y_j \rangle_{i \in I, j \in J} \in \Sigma_X$. In particular, $\{z_1, \dots, z_s, x_1, \dots, x_m\}$ is a part of a basis of the lattice N .

Proof of lemma 6. Suppose that $\tau + \langle y_1 \rangle \notin \Sigma_X$. Then, up to renumbering $\{z_1, \dots, z_s\}$, there is a primitive collection $\{z_j, \dots, z_s, y_1\}$ with $1 \leq j \leq s$, and by lemma 3 we get primitive relations:

$$z_j + \dots + z_s + y_1 - (u_1 + \dots + u_{s-j+1}) = 0, \quad u_1 + \dots + u_{s-j+1} + x_1 - (z_j + \dots + z_s) = 0.$$

Now, theorem 2 implies that either $\tau + \langle x_1 \rangle \in \Sigma_X$, or $\{z_1, \dots, z_s\} \supseteq \{u_1, \dots, u_{s-j+1}\}$. This last case is impossible, because it would give a relation of linear dependence among $\{z_1, \dots, z_s, y_1\}$, against the hypotheses. Therefore $\tau + \langle x_1 \rangle \in \Sigma_X$, and by recurrence we have the statement. $\blacksquare$

Useful Remark. Suppose that we have a relation among generators:

$$a_1 x_1 + \dots + a_h x_h - (b_1 y_1 + \dots + b_k y_k) = 0$$

with $a_i, b_j \in \mathbb{Z}_{>0}$ for each i, j . Then the two cones $\langle x_1, \dots, x_h \rangle$ and $\langle y_1, \dots, y_k \rangle$ can not be both in Σ_X , because they intersect in their interior, while two cones in the fan always intersect along a face.

Proof of theorem 5. We fix an m -dimensional linear subspace H' of H spanned by m pairs of centrally symmetric generators $x_i + y_i = 0$, with $x_1, \dots, x_m$ linearly independent.

Step 1: every generator of Σ_X contained in H' and different from $x_1, y_1, \dots, x_m, y_m$ has the form $\sum_{i=1}^m \varepsilon_i x_i$ with $\varepsilon_i \in \{1, 0, -1\}$ for all i .

Proof of step 1. Let $x_0 \in G(\Sigma_X)$ be a generator in H' different from $\{x_1, y_1, \dots, x_m, y_m\}$. Up to reorder $\{x_1, \dots, x_m\}$, we can suppose that $x_0 = \sum_{i=1}^s \varepsilon_i x_i$, with $2 \leq s \leq m$ and $\varepsilon_i \neq 0$ for all $i = 1, \dots, s$. Then for all $j \in \{1, \dots, s\}$ the generators $\{x_0, x_1, \dots, \check{x}_j, \dots, x_s\}$ are linearly independent, and by lemma 6 they are a part of a basis of the lattice, hence $|\varepsilon_j| = 1$. $\blacksquare$

We assume from now on that $x_0 = y_1 + \dots + y_m \in G(\Sigma_X)$.

Step 2: $m = 2r$ is even and Σ_X contains all the cones

$$\langle x_i, y_j \rangle_{i \in I^r, j \in J^r}, \quad \langle x_0, x_i, y_j \rangle_{i \in \tilde{I}^{r-1}, j \in \tilde{J}^r} \quad (\star)$$

and the primitive extremal relations

$$x_0 + \sum_{i \in I^r} x_i - \left(\sum_{j \in J^r} y_j \right) = 0, \quad \sum_{j \in J^{r+1}} y_j - (x_0 + \sum_{i \in I^{r-1}} x_i) = 0$$

for all partitions I^r, J^r and I^{r-1}, J^{r+1} of $\{1, \dots, m\}$ and for all choices of $\tilde{I}^{r-1}, \tilde{J}^r \subset \{1, \dots, m\}$ disjoint.

Moreover, for every $\tau \in \Sigma_X$ such that $\tau \cap H' = \{0\}$ and $\tau + \eta \in \Sigma_X$ for some η in $(\star)$, then $\tau + \eta' \in \Sigma_X$ for all η' in $(\star)$.

Proof of step 2. By lemma 6, since $\{x_0, x_1, \dots, x_{m-1}\}$ are linearly independent and $\langle x_0 \rangle \in \Sigma_X$, up to reorder $\{x_1, \dots, x_{m-1}\}$ we have $\langle x_0, x_1, \dots, x_{r-1}, y_r, \dots, y_{m-1} \rangle \in \Sigma_X$ with $1 \leq r \leq m$. Applying again lemma 6 to the cone $\sigma = \langle x_1, \dots, x_{r-1}, y_r, \dots, y_{m-1} \rangle$, we get that at least one of the two cones $\sigma + \langle y_m \rangle$, $\sigma + \langle x_m \rangle$ is in Σ_X .

Consider the relation $y_r + \dots + y_m - (x_0 + x_1 + \dots + x_{r-1}) = 0$. Since $\langle x_0, x_1, \dots, x_{r-1} \rangle \in \Sigma_X$, by lemma 1 this relation corresponds to an effective class, and since X is Fano, its degree must be positive: this gives $2r \leq m$. Moreover, it can not be $\langle y_r, \dots, y_m \rangle \in \Sigma_X$ (see the Useful Remark), hence $\sigma + \langle x_m \rangle = \langle x_0, x_1, \dots, x_{r-1}, y_r, \dots, y_{m-1}, x_m \rangle \in \Sigma_X$. This implies that also the relation $x_0 + x_1 + \dots + x_{r-1} + x_m - (y_r + \dots + y_{m-1}) = 0$ is effective, hence its degree is positive, and we get $2r \geq m$. Therefore $2r = m$. In particular, the relation $x_0 + x_1 + \dots + x_{r-1} + x_m - (y_r + \dots + y_{m-1}) = 0$ has degree 1, so by theorem 2 $\langle x_1, \dots, x_{r-1}, x_m, y_r, \dots, y_{m-1} \rangle \in \Sigma_X$.

Now, we know that at least one cone in each of the two types in $(\star)$ is in Σ_X . Let I^r , J^r be such that the cone $\langle x_i, y_j \rangle_{i \in I^r, j \in J^r}$ is in Σ_X . Then the relation

$$x_0 + \sum_{i \in I^r} x_i - \left(\sum_{j \in J^r} y_j \right) = 0$$

is effective by lemma 1, thus theorem 2 implies that the cones

$$\langle x_0, x_i, y_j \rangle_{i \in I^r \setminus \{p\}, j \in J^r}$$

are in Σ_X for all $p \in I^r$.

In an analogue way, suppose that for some $\tilde{I}^{r-1}$, $\tilde{J}^r$ the cone $\langle x_0, x_i, y_j \rangle_{i \in \tilde{I}^{r-1}, j \in \tilde{J}^r}$ is in Σ_X . Let $\{1, \dots, m\} = \tilde{I}^{r-1} \cup \tilde{J}^r \cup \{l\}$. Then the relations

$$\sum_{j \in \tilde{J}^r \cup \{l\}} y_j - (x_0 + \sum_{i \in \tilde{I}^{r-1}} x_i) = 0, \quad x_0 + \sum_{i \in \tilde{I}^{r-1} \cup \{l\}} x_i - \left(\sum_{j \in \tilde{J}^r} y_j \right) = 0$$

are effective by lemma 1, and again, theorem 2 implies that the cones

$$\langle x_0, x_i, y_j \rangle_{i \in \tilde{I}^{r-1}, j \in (\tilde{J}^r \setminus \{p\}) \cup \{l\}}, \quad \langle x_i, y_j \rangle_{i \in \tilde{I}^{r-1} \cup \{l\}, j \in \tilde{J}^r}$$

are in Σ_X for all $p \in \tilde{J}^r$. Hence, proceeding in this way, we get all the desired cones and extremal relations in Σ_X .

Finally, if τ is such that $\tau \cap H' = \{0\}$ and $\tau + \eta \in \Sigma_X$ for some η in $(\star)$, then it is easy to see, using the extremal relations and reasoning as above, that $\tau + \eta' \in \Sigma_X$ for all η' in $(\star)$. $\blacksquare$

Step 3: set $y_0 = -x_0$. Either $H' \cap G(\Sigma_X) = \{x_0, x_1, y_1, \dots, x_m, y_m\}$, or $H' \cap G(\Sigma_X) = \{x_0, y_0, x_1, y_1, \dots, x_m, y_m\}$.

Proof of step 3. Suppose that $z \in H' \cap G(\Sigma_X)$, $z \notin \{x_0, x_1, y_1, \dots, x_m, y_m\}$: we want to show that $z = y_0 = -x_0$. We know by step 1 that $z = \sum_{i=1}^m \lambda_i y_i$ with $\lambda_i \in \{1, 0, -1\}$ for all i :

$$z = \sum_{\lambda_i=1} y_i + \sum_{\lambda_i=-1} x_i = x_0 + \sum_{\lambda_i=0} x_i + \sum_{\lambda_i=-1} 2x_i.$$

Since z can not be in the interior of some cone of Σ_X , we know by step 2 that either $\#\{i \mid \lambda_i = 1\} \geq r + 1$ or $\#\{i \mid \lambda_i = -1\} \geq r + 1$, and that $\#\{i \mid \lambda_i = 0 \text{ or } \lambda_i = -1\} \geq r$. Hence at least $r + 1$ of the λ_i are -1. Thus we have

$$x_0 + z = \sum_{i=1}^m (\lambda_i + 1)y_i = \sum_{\lambda_i=1} 2y_i + \sum_{\lambda_i=0} y_i$$

and since there are at most $r - 1$ y_i appearing on the right hand side, we know by step 2 that they generate a cone in Σ_X . Hence $\{x_0, z\}$ is a primitive collection (see the Useful Remark), and since X is Fano, up to reordering $\{y_1, \dots, y_m\}$ there are only two possible primitive relations: either $x_0 + z = 0$ (so $z = y_0$), or $x_0 + z - y_1 = 0$ (so $z = x_2 + \dots + x_m$). Let's show that this last case is impossible. If $x_0 + z - y_1 = 0$, this relation has degree 1; thus by theorem 2 $\langle y_1, \dots, y_r \rangle \in \Sigma_X$ implies $\langle z, y_1, \dots, y_r \rangle \in \Sigma_X$. But we also have $z + y_2 + \dots + y_r = x_{r+1} + \dots + x_m$ and $\langle x_{r+1}, \dots, x_m \rangle \in \Sigma_X$, a contradiction (by the Useful Remark). ■

Step 4: for every generator $w \in G(\Sigma_X)$ such that $w \notin H'$, we have $\langle w \rangle + \eta \in \Sigma_X$ for all η in $(\star)$.

Proof of step 4. Suppose first that $\langle w, x_0 \rangle \in \Sigma_X$. Since $\{w, x_0, x_1, \dots, x_{m-1}\}$ are linearly independent, we know by lemma 6 that there exists a partition I, J of $\{1, \dots, m - 1\}$ such that $\langle w, x_0, x_i, y_j \rangle_{i \in I, j \in J} \in \Sigma_X$. This cone can not contain a primitive collection, so it must be $\#I \leq r - 1$, $\#J \leq r$. Since $\#I + \#J = 2r - 1$, it must be $\#I = r - 1$, $\#J = r$. Hence the statement follows from step 2.

If $\langle w, x_0 \rangle \notin \Sigma_X$, it means that $\{w, x_0\}$ is a primitive collection. Since w is not in H' , $w \neq -x_0$; hence the only possible primitive relation is $w + x_0 - u = 0$, which has degree 1 and is extremal by theorem 2. Now, $u \notin H'$ and $\langle u, x_0 \rangle \in \Sigma_X$: so by what precedes, $\langle u \rangle + \eta \in \Sigma_X$ for all η in $(\star)$. Choose $\eta = \langle x_i, y_j \rangle_{i \in I^r, j \in J^r}$ for some I^r, J^r . Since the relation $w + x_0 - u = 0$ is extremal, theorem 2 implies that also $\langle w \rangle + \eta$ is in Σ_X . By step 2, the same is true for all η' in $(\star)$, thus we get $\langle w, x_0 \rangle \in \Sigma_X$, a contradiction. ■

Step 5: let W be the invariant subvariety associated to one of the cones in $(\star)$. Then W is Fano.

Proof of step 5. We remark first of all that the linear span of the cones in $(\star)$ is always H' , and by step 2 the invariant subvarieties associated to these cones are all isomorphic. Let's fix a cone η in $(\star)$ and let $W = V(\eta)$.

Consider an invariant curve C in W : in Σ_X the curve C is associated to the $(n - 1)$ -dimensional cone $\langle w_3, \dots, w_s \rangle + \eta = (\langle w_1, w_3, \dots, w_s \rangle + \eta) \cap (\langle w_2, w_3, \dots, w_s \rangle + \eta)$. Remark that by step 2, $\langle w_1, w_3, \dots, w_s \rangle + \eta' \in \Sigma_X$ and $\langle w_2, w_3, \dots, w_s \rangle + \eta' \in \Sigma_X$ for all η' in $(\star)$. The numerical class of C in W is given by the relation

$$\sum_{i=1}^s a_i \bar{w}_i = 0,$$

where $a_i \in \mathbb{Z}$ and $\bar{w}$ denotes the image of $w \in N$ under the projection $N \rightarrow N/H'$. Hence in W the anticanonical degree of C is $\sum_i a_i$. In the lattice N , this relation lifts as

$$\sum_{i=1}^s a_i w_i + \sum_{j=1}^m \lambda_j x_j = 0,$$

where $\lambda_j \in \mathbb{Z}$. Choose a partition I^r, J^r of $\{1, \dots, m\}$: then the two relations

$$\sum_{i=1}^s a_i w_i + \sum_{k \in I^r} \lambda_k x_k - \sum_{j \in J^r} \lambda_j y_j = 0, \quad \sum_{i=1}^s a_i w_i + \sum_{j \in J^r} \lambda_j x_j - \sum_{k \in I^r} \lambda_k y_k = 0$$

correspond respectively to the numerical classes of the invariant curves in X :

$$V(\langle w_3, \dots, w_s, x_k, y_j \rangle_{k \in I^r, j \in J^r}), \quad V(\langle w_3, \dots, w_s, x_j, y_k \rangle_{j \in J^r, k \in I^r}).$$

Since X is Fano, they both have positive degree, so $\sum_i a_i \geq 1 + |\sum_{k \in I^r} \lambda_k - \sum_{j \in J^r} \lambda_j| \geq 1$. $\blacksquare$

Step 6: Suppose that $y_0 = -x_0 \in H' \cap G(\Sigma_X)$. Then every cone $\sigma \in \Sigma_X$ decomposes as $\sigma = \eta + \tau$, with $\eta \subset H'$, $\dim \eta = m$ and $\tau \cap H' = \{0\}$; H' contains the fan of a Del Pezzo variety V^m and X is a toric bundle over W with fiber V^m .

Proof of step 6. Apply step 2 twice considering first x_0 and then y_0 : it is immediate to see that H' contains the fan of the Del Pezzo variety V^m . Consider now a cone $\tau \in \Sigma_X$ such that $\tau \cap H' = \{0\}$. By lemma 6, we know that $\tau + \eta \in \Sigma_X$ for at least one m -dimensional $\eta \subset H'$. Hence, again by step 2, $\tau + \eta' \in \Sigma_X$ for all $\eta' \subset H'$. Thus by theorem 4 X is a toric bundle over a smooth toric variety with fiber V^m . Since W is an invariant section of the bundle, it is isomorphic to the basis of the bundle. $\blacksquare$

Step 7: suppose that $H' \cap G(\Sigma_X) = \{x_0, x_1, y_1, \dots, x_m, y_m\}$. Then there exist a smooth projective toric variety X' with $G(\Sigma_{X'}) = G(\Sigma_X)$, such that X' is a toric bundle over W with fiber the pseudo Del Pezzo variety $\tilde{V}^m$.

Proof of step 7. We remark that by step 4, all generators in $G(\Sigma_X) \setminus H'$ induce a generator in Σ_W . The fan $\Sigma_{X'}$ is defined as follows:

1. $G(\Sigma_{X'}) = G(\Sigma_X)$;
2. the set $\{\sigma \in \Sigma_{X'} \mid \sigma \subset H'\}$ is the fan of the pseudo Del Pezzo variety $\tilde{V}^m$ in H' , with generators $\{x_0, x_1, y_1, \dots, x_m, y_m\}$;
3. for every $\tau \in \Sigma_X$ such that $\tau \cap H' = \{0\}$ and $\tau + \eta \in \Sigma_X$ for η in $(\star)$, and for every $\sigma \subset H'$ in the fan of $\tilde{V}^m$, we set $\tau + \sigma \in \Sigma_{X'}$.

By construction, X' is a toric bundle with fiber $\tilde{V}^m$ (see theorem 4). The invariant subvarieties $V(\sigma) \subset X'$ for $\sigma \in \Sigma_{X'}(m)$, $\sigma \subset H'$, are invariant sections of the bundle. If σ is one of the cones in $(\star)$, then $X \dashrightarrow X'$ is an isomorphism on $V(\sigma) \simeq W$. Therefore the basis of the bundle is actually W . $\blacksquare$

Let's consider now the linear subspace H of $N_{\mathbb{Q}}$ spanned by all pairs of symmetric generators in Σ_X . We want to prove that H decomposes as a direct sum of subspaces as H' .

Fix $h = \dim H$ pairs $x_i + y_i = 0$ that span H . If $G(\Sigma_X) \cap H = \{x_1, y_1, \dots, x_h, y_h\}$, then the desired decomposition is just $H = K$. If there is a generator $x_0 \in H$ different from $\{x_i, y_i\}$, by step 1, up to renaming $\{x_i, y_i\}$, we can suppose that $x_0 = y_1 + \dots + y_m$ with $2 \leq m \leq h$. Let H' be the linear span of $\{x_1, \dots, x_m\}$. We have to show that if v is another generator in H such that $v \neq -x_0$, then the smallest linear subspace H'' containing v satisfies $H' \cap H'' = \{0\}$.

Again by step 1, we know that $v = \sum_{i=1}^h \varepsilon_i x_i$ with $\varepsilon_i \in \{1, 0, -1\}$ for all i . Up to renaming $\{x_{m+1}, y_{m+1}, \dots, x_h, y_h\}$, we can assume that

$$v = \sum_{i \in I} x_i + \sum_{j \in J} y_j + \sum_{k=m+1}^s y_k$$

where $I, J \subset \{1, \dots, m\}$ disjoint and $m < s \leq h$. Remark that it must be $s > m$, because by step 3 we know that $v \notin H'$ (and $x_0 \notin H''$).

Suppose first that $\#I \leq r$ and $\#J \leq r$ and choose a partition I^r, J^r of $\{1, \dots, m\}$ such that $I^r \supseteq I$, $J^r \supseteq J$. Set $\eta = \langle x_i, y_j \rangle_{i \in I^r, j \in J^r}$. By step 4, we have $\langle v \rangle + \eta \in \Sigma_X$, and since $\{v, x_i, y_j, x_{m+1}, \dots, x_{s-1}\}_{i \in I^r, j \in J^r}$ are linearly independent, lemma 6 implies that we can reorder $\{x_{m+1}, \dots, x_{s-1}\}$ in such a way that

$$\eta + \langle v, x_{m+1}, \dots, x_t, y_{t+1}, \dots, y_{s-1} \rangle \in \Sigma_X, \text{ with } m \leq t \leq s-1,$$

and by step 2 the same is true for $\eta' = \langle x_j, y_i \rangle_{j \in J^r, i \in I^r}$. Now, the relations

$$\sum_{k=t+1}^s y_k - (v + \sum_{i \in I} y_i + \sum_{j \in J} x_j + \sum_{h=m+1}^t x_h) = 0, \quad v + \sum_{h=m+1}^t x_h + x_s - (\sum_{i \in I} x_i + \sum_{j \in J} y_j + \sum_{k=t+1}^{s-1} y_k) = 0$$

are effective by lemma 1, and since X is Fano, they have positive degree: thus we get $\#I + \#J \leq s + m - 2t - 2$ and $\#I + \#J \leq 2t + 2 - m - s$. So $\#I + \#J \leq 0$ and $\#I = \#J = 0$: then $v = \sum_{k=m+1}^s y_k$ and $H' \cap H'' = \{0\}$.

Suppose now that either $\#I$ or $\#J$ is bigger than r . In this case, we get a contradiction; the proof is very similar to the preceeding case. Suppose that $\#I > r$: we can write

$$v = x_1 + \dots + x_r + \sum_{i \in I'} x_i + \sum_{j \in J} y_j + \sum_{k=m+1}^s y_k,$$

where $I = \{1, \dots, r\} \cup I'$. Reasonning as before we get the same cones in Σ_X ; now we consider the relations:

$$\begin{aligned} \sum_{i \in I'} x_i + \sum_{k=t+1}^s y_k - (v + y_1 + \dots + y_r + \sum_{j \in J} x_j + \sum_{h=m+1}^t x_h) &= 0, \\ v + \sum_{i \in I'} y_i + \sum_{h=m+1}^t x_h + x_s - (x_1 + \dots + x_r + \sum_{j \in J} y_j + \sum_{k=t+1}^{s-1} y_k) &= 0. \end{aligned}$$

Since they have positive degree, we get $\#I' \geq \#J + r$, hence $\#J = 0$ and $\#I' = r$. Thus $v = -x_0$, a contradiction. In the same way, if $\#J > r$, we get $v = x_0$, again a contradiction. $\blacksquare$

Remark. In the same setting as step 7, it is not true in general that H' contains the fan of $\tilde{V}^m$. Indeed, the sets $\{x_i, y_j\}_{i \in I^{r+s}, j \in J^{r-s}}$ where I^{r+s}, J^{r-s} is a partition of $\{1, \dots, m\}$ and $s \in \{1, \dots, r\}$, do not need to generate a cone in Σ_X .

As an immediate corollary, we get a generalization of Ewald's result [Ewa88]:

Proposition 7. *Let X be a toric Fano variety and let H be the linear subspace of $N_{\mathbb{Q}}$ spanned by all pairs of symmetric generators in $G(\Sigma_X)$.*

Suppose that $\dim H = n$. Then X is a product of Del Pezzo varieties, pseudo Del Pezzo varieties and a power of $\mathbb{P}^1$.

Suppose that $\dim H = n - 1$. Then there exists $X' \in \mathcal{B}_X$ which is a toric bundle over $\mathbb{P}^1$ with fiber a product of Del Pezzo varieties, pseudo Del Pezzo varieties and a power of $\mathbb{P}^1$.

In dimension 4, there are only two toric Fano varieties which have 3 linearly independent pairs of symmetric generators and such that X' is not isomorphic to X . These 4-folds are $M1$ and $R3$, in Batyrev's notation [Bat99]. In both cases $\varphi: X' \dashrightarrow X$ is just a flip.

Proof of proposition 7. The statement for $\dim H = n$ is straightforward from theorem 5: let $X \dashrightarrow X' \rightarrow W$ be as in theorem 5, (iii). Then $G(\Sigma_W)$ consists of $\dim W$ linearly independent pairs of symmetric generators $\bar{x}_i, \bar{y}_i$, where $\bar{u}$ denotes the image of u under the projection $N \rightarrow N/H'$, $H' = H_1 \oplus \dots \oplus H_a \oplus \tilde{H}_1 \oplus \dots \oplus \tilde{H}_b$. Since W is Fano, we have $W \simeq (\mathbb{P}^1)^{\dim W}$. Moreover, the primitive relations of W $\bar{x}_i + \bar{y}_i = 0$ remain unchanged when lifted to N , hence X' is actually the product $V^{h_1} \times \dots \times V^{h_a} \times \tilde{V}^{\tilde{h}_1} \times \dots \times \tilde{V}^{\tilde{h}_b} \times (\mathbb{P}^1)^{\dim W}$. Since X' is Fano, $X' \simeq X$.

Assume $\dim H = n - 1$ and fix $n - 1$ pairs x_i, y_i with $i = 1, \dots, n - 1$ such that $x_1, \dots, x_{n-1}$ are a basis for H and $x_i + y_i = 0$ for all i .

Claim. Let $v, w \in G(\Sigma_X) \setminus H$. Then either $\{v, w\}$ is a primitive collection with relation $v + w - z = 0$ and $z \in G(\Sigma_X) \cap H$, or $\langle v, w \rangle \in \Sigma_X$ and $w = \sum_{i=1}^{n-1} \varepsilon_i x_i + \delta v$ with $|\delta| = 1$ and $\varepsilon_i \in \{1, 0, -1\}$ for all i ; moreover, the number of non-zero ε_i is odd.

Proof of the claim. If $v \in G(\Sigma_X) \setminus H$, then by lemma 6 $\{v, x_1, \dots, x_{n-1}\}$ is a basis of N . Hence, if $v, w \in G(\Sigma_X) \setminus H$, we have $w = \sum_{i=1}^{n-1} \varepsilon_i x_i + \delta v$ with $|\delta| = 1$.

Suppose that $\{v, w\}$ is a primitive collection. Since $v + w \neq 0$ (otherwise $\dim H = n$), the associated primitive relation must be $v + w - z = 0$, with $z \in G(\Sigma_X)$. Then $z = \sum_{i=1}^{n-1} \varepsilon_i x_i + (1 + \delta)v$, hence it must be $\delta = -1$ and $z \in H$.

Let's assume that $\langle v, w \rangle \in \Sigma_X$. We can reorder $\{x_1, \dots, x_{n-1}\}$ in such a way that $w = \sum_{i=1}^r \varepsilon_i x_i + \delta v$ with $r \in \{1, \dots, n-1\}$ and $\varepsilon_i \neq 0$ for all $i = 1, \dots, r$. Fix $i \in \{1, \dots, r\}$. Since $\{v, w, x_1, \dots, \tilde{x}_i, \dots, x_r\}$ are linearly independent, lemma 6 implies that it is a part of a basis of the lattice. Hence $|\varepsilon_i| = 1$.

It remains to show that r is odd. Up to renaming $\{x_i, y_i\}$, we can assume that $w = \sum_{i=1}^r x_i + \delta v$, where $|\delta| = 1$. Again by lemma 6, there exists a partition I, J of $\{1, \dots, r-1\}$ such that $\langle v, w, x_i, y_j \rangle_{i \in I, j \in J} \in \Sigma_X$. Consider now the two relations:

$$\sum_{i \in I} x_i + x_r + \delta v - w - \sum_{j \in J} y_j = 0, \quad \sum_{j \in J} y_j + y_r + w - \delta v - \sum_{i \in I} x_i = 0.$$

They are both effective by lemma 1, so they must have positive degree: this gives $\#J = \#I + \delta - 1$, hence $r = 2\#I + \delta$ is odd. $\blacksquare$

We first prove the proposition in the case $G(\Sigma_X) \cap H = \{x_1, y_1, \dots, x_{n-1}, y_{n-1}\}$ (namely, in the decomposition given by theorem 5, we have $H = K$). Then, we can restate the claim as follows: for every $v, w \in G(\Sigma_X) \setminus H$, we have $w = \sum_{i=1}^{n-1} \varepsilon_i x_i + \delta v$ with $|\delta| = 1$ and $\varepsilon_i \in \{1, 0, -1\}$ for all i ; moreover, the number of non-zero ε_i is odd.

Remark that $\#(G(\Sigma_X) \setminus H) \geq 2$, because since X is complete, there is at least one generator in both half-spaces cut by H . Let's show that $\#(G(\Sigma_X) \setminus H) = 2$. By contradiction, suppose that $G(\Sigma_X) \setminus H \supseteq \{v_1, v_2, w\}$ with v_1, v_2 in the same half-space cut by H and w in the other one. By the claim we have $v_2 = v_1 + \sum_{i \in I} \varepsilon_i x_i$ and $w = -v_1 + \sum_{j \in J} \delta_j x_j$ with $I, J \subseteq \{1, \dots, n-1\}$, $|\varepsilon_i| = |\delta_j| = 1$ for all i, j , and $\#I$ and $\#J$ odd. Then

$$w + v_2 = \sum_{i \in (I \setminus J)} \varepsilon_i x_i + \sum_{j \in (J \setminus I)} \delta_j x_j + \sum_{i \in I \cap J} (\varepsilon_i + \delta_i) x_i.$$

Since by the claim all the coefficients must be 1 or -1, we have $\varepsilon_i + \delta_i = 0$ for all $i \in I \cap J$, and the non-zero coefficients are $\#(I \setminus J) + \#(J \setminus I) = \#I + \#J - 2\#(I \cap J)$ an even number, a contradiction.

Therefore we have $G(\Sigma_X) = \{x_1, y_1, \dots, x_{n-1}, y_{n-1}, v, w\}$ and up to renaming $\{x_i, y_i\}$, we can assume $v + w = x_1 + \dots + x_r$, $r \in \{1, \dots, n-1\}$ odd. Hence $G(\Sigma_X) = G(\Sigma_{X'})$ where X' has primitive relations: $x_i + y_i = 0$ for $i = 1, \dots, n-1$, and $v + w - (x_1 + \dots + x_r) = 0$. Thus X' is a $(\mathbb{P}^1)^{n-1}$ -bundle over $\mathbb{P}^1$. We can actually say more: define Y' to be the $(r+1)$ -dimensional smooth toric variety with $G(\Sigma_{Y'}) = \{x_1, y_1, \dots, x_r, y_r, v, w\}$ and primitive relations $x_i + y_i = 0$ for $i = 1, \dots, r$ and $v + w - (x_1 + \dots + x_r) = 0$. Then Y' is a $(\mathbb{P}^1)^r$ -bundle over $\mathbb{P}^1$ and $X' = (\mathbb{P}^1)^{n-1-r} \times Y'$. Moreover, $X = (\mathbb{P}^1)^{n-1-r} \times Y$, where Y is Fano and $G(\Sigma_Y) = G(\Sigma_{Y'})$.

We have now to prove the general case. Let $X \dashrightarrow X' \rightarrow W$ be as in theorem 5, (iii). Then W is Fano and has at least $\dim W - 1$ linearly independent pairs of symmetric generators $\bar{x}_i + \bar{y}_i = 0$, where $\bar{u}$ denotes the image of u under the projection $N \rightarrow N/H'$, $H' = H_1 \oplus \dots \oplus H_a \oplus \tilde{H}_1 \oplus \dots \oplus \tilde{H}_b$. By what precedes, $G(\Sigma_W)$ has exactly $2 \dim W$ elements: the last two are $\bar{v}, \bar{w}$ such that $\bar{v} + \bar{w} = \bar{x}_1 + \dots + \bar{x}_r$ for some $r \in \{0, \dots, \dim W\}$ (if $r > 0$ then it must be odd). In X' , by hypothesis we still have $x_i + y_i = 0$; the last relation lifts as $v + w = x_1 + \dots + x_r + u_1 + \dots + u_s$, where $\{u_1, \dots, u_s\}$ are contained in H' (and again $r + s$ is odd). Hence X' is a toric bundle over $\mathbb{P}^1$ with fiber $V^{h_1} \times \dots \times V^{h_a} \times \tilde{V}^{\tilde{h}_1} \times \dots \times \tilde{V}^{\tilde{h}_b} \times (\mathbb{P}^1)^{\dim W - 1}$. $\blacksquare$

Remark. We think that it should be possible, with a detailed analysis, to classify all the possible X Fano with $\dim H = n - 1$, namely: to list all the possible decompositions of

$v + w$ in H , or in other words again, to list all the possible $X' \rightarrow \mathbb{P}^1$ with fiber a product of Del Pezzo varieties, pseudo Del Pezzo varieties and a power of $\mathbb{P}^1$, such that there exists X Fano with $G(\Sigma_X) = G(\Sigma_{X'})$. In the proof, we did this when the fiber of $X' \rightarrow \mathbb{P}^1$ is just a power of $\mathbb{P}^1$.

Acknowledgements. I would like to thank Laurent Bonavero and Olivier Debarre, for many useful conversations about toric Fano varieties.

I am also very grateful to Lucia Caporaso, my advisor, for her constant guide.

References

- [Bat82] Victor V. Batyrev. Toroidal Fano 3-folds. *Mathematics of the USSR Izvestiya*, 19:13–25, 1982.
- [Bat99] Victor V. Batyrev. On the classification of toric Fano 4-folds. *Journal of Mathematical Sciences (New York)*, 94:1021–1050, 1999.
- [BCDD02] Laurent Bonavero, Cinzia Casagrande, Olivier Debarre, and Stéphane Druel. Sur une conjecture de Mukai. Prépublication 566, Institut Fourier, 2002. Preprint math.AG/0204314.
- [BCW02] Laurent Bonavero, Frédéric Campana, and Jaroslaw A. Wiśniewski. Variétés projectives complexes dont l'éclatée en un point est de Fano. *Comptes Rendus de l'Académie des Sciences*, 334:463–468, 2002.
- [Bon00] Laurent Bonavero. Toric varieties whose blow-up at a point is Fano. Preprint math.AG/0012229, 2000. To appear in Tohoku Mathematical Journal.
- [Cas01a] Cinzia Casagrande. Contractible classes in toric varieties. Preprint math.AG/0111332, 2001. To appear in Mathematische Zeitschrift.
- [Cas01b] Cinzia Casagrande. On the birational geometry of toric Fano 4-folds. *Comptes Rendus de l'Académie des Sciences*, 332:1093–1098, 2001.
- [Cas01c] Cinzia Casagrande. Toric Fano varieties and birational morphisms. Preprint math.AG/0112007, 2001.
- [Ewa88] Günter Ewald. On the classification of toric Fano varieties. *Discrete Computational Geometry*, 3:49–54, 1988.
- [Ewa96] Günter Ewald. *Combinatorial Convexity and Algebraic Geometry*, volume 168 of *Graduate Texts in Mathematics*. Springer-Verlag, 1996.
- [Ful93] William Fulton. *Introduction to Toric Varieties*. Number 131 in Annals of Mathematics Studies. Princeton University Press, 1993.
- [Oda88] Tadao Oda. *Convex Bodies and Algebraic Geometry - An Introduction to the Theory of Toric Varieties*. Springer-Verlag, 1988.
- [Rei83] Miles Reid. Decomposition of toric morphisms. In *Arithmetic and Geometry, vol. II: Geometry*, number 36 in Progress in Mathematics, pages 395–418. Birkhäuser, 1983.
- [Sat00] Hiroshi Sato. Toward the classification of higher-dimensional toric Fano varieties. *Tohoku Mathematical Journal*, 52:383–413, 2000.

- [Sat02] Hiroshi Sato. Toric Fano varieties with divisorial contractions to curves. Preprint math.AG/0205298, 2002.
- [VK85] V. E. Voskresenskii and Alexander Klyachko. Toroidal Fano varieties and roots systems. *Mathematics of the USSR Izvestiya*, 24:221–244, 1985.
- [Wiś02] Jarosław A. Wiśniewski. Toric Mori theory and Fano manifolds. In *Geometry of Toric Varieties*, volume 6 of *Séminaires et Congrès*, pages 249–272. Société Mathématique de France, 2002.
- [WW82] Keiichi Watanabe and Masayuki Watanabe. The classification of Fano 3-folds with torus embeddings. *Tokyo Journal of Mathematics*, 5:37–48, 1982.